

DIMENSION PROPERTIES OF THE BOUNDARIES OF EXPONENTIAL BASINS

KRZYSZTOF BARAŃSKI, BOGUSŁAWA KARPIŃSKA, AND ANNA ZDUNIK

ABSTRACT. We prove that the boundary of a component U of the basin of an attracting periodic cycle (of period greater than 1) for an exponential map on the complex plane has Hausdorff dimension greater than 1 and less than 2. Moreover, the set of points in the boundary of U which do not escape to infinity has Hausdorff dimension (in fact: hyperbolic dimension) greater than 1, while the set of points in the boundary of U which escape to infinity has Hausdorff dimension 1.

1. INTRODUCTION

Let

$$f(z) = \lambda \exp(z)$$

for $\lambda \in \mathbb{C} \setminus \{0\}$, $z \in \mathbb{C}$ be an exponential map with an attracting periodic cycle of period p . Then the Julia set $J(f)$ is equal to the boundary of the (entire) basin B of attraction to this cycle. We are interested in dimension properties of this boundary. We consider also the escaping set

$$I(f) = \{z \in \mathbb{C} : f^n(z) \rightarrow \infty \text{ as } n \rightarrow \infty\}$$

and the set of points in the Julia set with bounded forward trajectories

$$J_{\text{bd}}(f) = \{z \in J(f) : \{f^n(z)\}_{n=0}^{\infty} \text{ is bounded}\}.$$

In our context, both set are contained in the boundary of B .

In the well-known case $p = 1$, the attracting basin consists of a unique simply connected component U . As proved in [11], the Hausdorff dimension $\dim_{\text{H}}$ of its boundary $\partial U = J(f)$ is then equal to 2. In fact, it follows from the proof in [11] that $\dim_{\text{H}}(\partial U) = \dim_{\text{H}}(\partial U \cap I(f)) = 2$. On the other hand, we have $1 < \dim_{\text{H}}(\partial U \cap J_{\text{bd}}(f)) \leq \dim_{\text{H}}(\partial U \setminus I(f)) < 2$ (see [8, 15]). A refined analysis of the dimension of subsets of $I(f)$ according to the speed of convergence to ∞ can be found in [10]. Hence, it follows from the above results that in the case $p = 1$ the dimension of ∂U is carried by the set of escaping points.

Consider now the case $p > 1$. Then the basin B of the attracting cycle consists of infinitely many disjoint simply connected components. Again, the Hausdorff dimension of the boundary of B is equal to 2 (see [11]). However, this does not imply that the

Date: February 6, 2009.

2000 Mathematics Subject Classification. Primary 37F10, 37F35, 30D40, 28A80.

Research supported by Polish MNiSW Grant N N201 0234 33 and EU FP6 Marie Curie Programmes RTN CODY and ToK SPADE2. The second author is supported by Polish PW Grant 504G 1120 0011 000.

boundaries of components of B have Hausdorff dimension 2. In fact, in this paper we prove the following theorem.

Theorem A. *Let $p > 1$ and let U be a component of the basin of the attracting cycle for the map f . Then the boundary of U has Hausdorff dimension greater than 1 and less than 2.*

This shows that for $p > 1$ the dimension of ∂U is carried by points in the Julia set, which are not in the boundary of any component of the Fatou set (so-called buried points). By Theorem A, we immediately obtain:

Corollary A'. *If $p > 1$, then the set of buried points for the map f has Hausdorff dimension 2.*

Surprisingly, for $p > 1$ the role of escaping and non-escaping points in ∂U from dimensional point of view is quite different than in the case $p = 1$. This is shown in the second result proved in this paper.

Theorem B. *If $p > 1$ and U is a component of the basin of the attracting cycle for the map f , then*

$$1 < \dim_{\mathrm{H}}(\partial U \cap J_{\mathrm{bd}}(f)) \leq \dim_{\mathrm{H}}(\partial U \setminus I(f)) = \dim_{\mathrm{H}}(\partial U) < 2,$$

while

$$\dim_{\mathrm{H}}(\partial U \cap I(f)) = 1.$$

Remark. In fact, the proof of Theorem B shows that $\partial U \cap J_{\mathrm{bd}}(f)$ has hyperbolic dimension greater than 1.

Therefore, in the case $p > 1$ the dimension of ∂U is carried by the set of non-escaping points.

Remark. It is easy to check that for $z \in \partial U$ we have

$$f^n(z) \xrightarrow{n \rightarrow \infty} \infty \iff f^{np}(z) \xrightarrow{n \rightarrow \infty} \infty$$

(see Lemma 7).

Recall that $J(f)$ consists of so-called hairs homeomorphic to $[0, \infty)$, tending to ∞ , composed of points with given symbolic itineraries. All points from a hair outside its endpoint (the point corresponding to 0 in $[0, \infty)$) are contained in $I(f)$. In the case $p = 1$ the hairs are pairwise disjoint and the Julia set is homeomorphic to a so-called straight brush (see [1]), while in the case $p > 1$ some hairs have common endpoints and the Julia set is a modified straight brush (see [5]). In both cases the Hausdorff dimension of the union of hairs without endpoints is equal to 1 (see [9, 13]). Hence, the hairs without endpoints are insignificant from dimensional point of view.

The plan of the paper is as follows. In Section 2 we describe combinatorics of exponential maps with a periodic basin of attraction and provide a symbolic description of the boundaries of its components (Proposition 3). Then, in Section 3 we prove that the set of escaping points in the boundary U has Hausdorff dimension 1. In Section 4 it is shown that

the set of points in the boundary of U with bounded trajectories has hyperbolic dimension greater than 1. Finally, Section 5 concludes the proof.

2. COMBINATORICS OF EXPONENTIALS WITH ATTRACTING PERIODIC BASINS

The combinatorics of the Julia sets of exponential maps with an attracting periodic cycle of period $p > 1$ was described in [4, 5]. For completeness, we briefly recall the construction from these papers.

Let $z_0, \dots, z_p = z_0$ be the attracting cycle and let $U_0, \dots, U_p = U_0$ be the components of the basin of attraction, such that $z_j \in U_j$. Of course, we have $f^p(U_j) \subset U_j$. All U_j are simply connected (see e.g. [2]). Since 0 is the only singular value of f , it must be in a component containing some z_j , so we can assume $0 \in U_1$. It is easy to find a simply connected domain $B_{p+1} \subset U_1$ such that $0, z_1 \in B_{p+1}$, ∂B_{p+1} is a Jordan curve in U_1 and $f^p(\overline{B_{p+1}}) \subset B_{p+1}$. Let $B_p = f^{-1}(B_{p+1})$. Then B_p is simply connected, contains some left-half plane and ∂B_p is a curve homeomorphic to the real line, contained in $\{z \in \mathbb{C} : a < \operatorname{Re}(z) < b\}$ for some $a, b \in \mathbb{R}$. This curve is periodic of period $2\pi i$ and its two “ends” have imaginary parts tending respectively to $\pm\infty$.

For $j = p-1, \dots, 0$ define successively B_j to be the component of $f^{-1}(B_{j+1})$ containing z_j . The construction implies that f maps univalently B_j onto B_{j+1} for $j = 1, \dots, p-1$, such that $B_1 \supset \overline{B_{p+1}}$, and f on $B_0 = f^{-1}(B_1)$ is a universal covering of $B_1 \setminus \{0\}$.

Let $C_p \subset B_p$ be a domain containing z_0 and some left half-plane $\{z : \operatorname{Re}(z) < c\}$, such that ∂C_p is a curve homeomorphic to the real line, its two “ends” have imaginary parts tending to $\pm\infty$, moreover ∂C_p coincides with ∂B_p on $\{z : |\operatorname{Im}(z)| < L_1\}$ and is vertical on $\{z : |\operatorname{Im}(z)| > L_2\}$ for some large constants $L_1, L_2 > 0$.

Define successively $C_j \subset B_j$ to be the component of $f^{-1}(C_{j+1})$ containing z_j for $j = p-1, \dots, 0$. Again, f maps univalently C_j onto C_{j+1} and f on $C_0 = f^{-1}(C_1)$ is a universal covering of $C_1 \setminus \{0\}$. Moreover, since $\overline{B_{p+1}}$ is compact, we have $\overline{B_{p+1}} \subset C_1$ if the constant L_1 is chosen sufficiently large. This implies

$$\overline{C_j} \subset U_j \quad \text{for } j = 0, \dots, p-1 \quad \text{and} \quad \overline{C_p} \subset C_0.$$

In particular, $\overline{C_0}, \dots, \overline{C_{p-1}}$ are pairwise disjoint. Since C_0 contains some left half-plane and $0 \notin C_0$, the set $\mathbb{C} \setminus C_0$ consists of infinitely many disjoint simply connected, closed, connected sets $\mathcal{H}_s$, $s \in \mathbb{Z}$, such that


$$(1) \quad \mathcal{H}_s = \mathcal{H}_0 + 2\pi i s$$

Similarly, for $j = 1, \dots, p-1$ the set $\overline{C_j}$ is a simply connected, closed, connected subset of $\bigcup_s \mathcal{H}_s$, such that

$$(2) \quad z_j \in C_j \subset U_j \subset \mathcal{H}_{k_j} \quad \text{for } j = 1, \dots, p-1,$$

for some $k_j \in \mathbb{Z}$. See Figure 1.

The following lemma is essentially stated as Proposition 2.10 in [4].

FIGURE 1. The sets C_j .

Lemma 1. For $s \in \mathbb{Z}$, $j = 1, \dots, p-1$, we have $\partial\mathcal{H}_s = \Gamma_s$, $\partial C_j = \gamma_j$, where $\Gamma_s, \gamma_j : \mathbb{R} \rightarrow \mathbb{C}$ are curves homeomorphic to the real line, such that

$$\operatorname{Re}(\Gamma_s(t)), \operatorname{Re}(\gamma_j(t)) \xrightarrow[t \rightarrow \pm\infty]{} +\infty.$$

Moreover, for sufficiently large $M > 0$, in the right half-plane $\{z : \operatorname{Re}(z) \geq M\}$ the curves Γ_s, γ_j form pairs of the graphs of smooth functions $y = H_s^\pm(x)$, $y = h_j^\pm(x)$, $x \geq M$, where (x, y) are Euclidean coordinates on the plane and

$$\begin{aligned} \lim_{x \rightarrow +\infty} H_s^\pm(x) &= (2s + 1 \pm 1)\pi - \operatorname{Arg}(\lambda), \\ \lim_{x \rightarrow +\infty} h_{p-1}^\pm(x) &= (2k_{p-1} + 1 \pm 1/2)\pi - \operatorname{Arg}(\lambda), \\ \lim_{x \rightarrow +\infty} h_j^\pm(x) &= 2\pi k_j - \operatorname{Arg}(\lambda) \quad \text{for } j = 1, \dots, p-2 \end{aligned}$$

for some branch of argument. Consequently,

$$\mathcal{H}_s \cap \{z : \operatorname{Re}(z) \geq M\} \subset \{z : (2s - 1)\pi - \operatorname{Arg}(\lambda) < \operatorname{Im}(z) < (2s + 3)\pi - \operatorname{Arg}(\lambda)\}.$$

Proof. By construction, $\gamma_{p-1} \cap \{z : \operatorname{Re}(z) \geq M\}$ for large M is a pair of curves, which are images of vertical half-lines under some branch of logarithm. Hence, these curves are the graphs of some smooth functions $h_{p-1}^\pm(x)$, such that $\lim_{x \rightarrow +\infty} h_{p-1}^\pm(x) = (2k_{p-1} + 1 \pm 1/2)\pi -$

$\text{Arg}(\lambda)$ and $|(h_{p-1}^\pm)'| < \varepsilon$ for a small $\varepsilon > 0$. Moreover, the curves γ_j , $j = p-2, \dots, 1$ and Γ_s are successive images of γ_{p-1} under inverse branches of f , which have the form $\log(z/\lambda)$ for some branch of logarithm. Note also that $|\text{Arg}(z)|, |\text{Arg}(\log'(z/\lambda))| < \varepsilon$ for every $z \in \bigcup_{j=1}^{p-1} \gamma_j$, provided $\text{Re}(z) \geq M$ for large M . This easily implies that γ_j and Γ_s on some right-half plane are pairs of smooth curves with almost horizontal tangent vector and the required asymptotics. $\square$

Since $\mathbb{C} \setminus C_1$ is simply connected and does not contain 0, we can define branches of f^{-1} ,

$$g_s : \mathbb{C} \setminus C_1 \rightarrow \mathcal{H}_s, \quad s \in \mathbb{Z},$$

mapping $\mathbb{C} \setminus C_1$ onto $\mathcal{H}_s$. Let

$$\mathcal{L} = \mathbb{C} \setminus \bigcup_{j=1}^p C_j$$

By construction, we have $\mathcal{L} \subset f(\mathcal{L})$, so

$$g_s(\mathcal{L}) \subset \mathcal{L}.$$

Hence, the collection of maps $\{g_s|_{\mathcal{L}}\}_{s \in \mathbb{Z}}$ forms an infinite conformal iterated function system.

Since the Julia set is the complement of the basin of attraction, for each $z \in J(f)$ all its forward iterates are contained in $\mathcal{L} \setminus C_0 \subset \bigcup_s \mathcal{H}_s$, so following [4], we can define its itinerary $(s_0, s_1, \dots)$, taking $s_n \in \mathbb{Z}$ such that $f^n(z) \in \mathcal{H}_{s_n}$, $n = 0, 1, \dots$. It is clear that

$$J(f) = \bigcap_{n=0}^{\infty} \bigcup_{s_0, \dots, s_n} g_{s_0} \circ \dots \circ g_{s_n}(\mathcal{L}),$$

and $\bigcap_{n=0}^{\infty} g_{s_0} \circ \dots \circ g_{s_n}(\mathcal{L})$ is precisely the set of points in the Julia set with itinerary $(s_0, s_1, \dots)$.

Let

$$\underline{k} = (k_1, \dots, k_{p-1})$$

be the block of $p-1$ symbols, where $k_1, \dots, k_{p-1}$ are integers defined in (2). See Figure 1. The block $\underline{k}$ describes the trajectory of the singular value 0 and is called the kneading sequence for f (see [4]). We will write

$$G_{\underline{k}} = g_{k_1} \circ \dots \circ g_{k_{p-1}}, \quad G_{s, \underline{k}} = g_s \circ g_{k_1} \circ \dots \circ g_{k_{p-1}}, \quad G_{\underline{k}, s} = g_{k_1} \circ \dots \circ g_{k_{p-1}} \circ g_s$$

for $s \in \mathbb{Z}$.

The following lemma is a kind of folklore (it follows from estimations of hyperbolic metric, see e.g. the proofs of Proposition 6.1 in [11] and Proposition 2.1 in [14]).

Lemma 2. *There exist $c > 0$, $A > 1$, such that for every $n > 0$, $s_0, \dots, s_n \in \mathbb{Z}$ and $z \in \mathcal{L}$,*

$$|(g_{s_0} \circ \dots \circ g_{s_n})'(z)| < cA^{-n}.$$

$\square$

Now we show how the boundary of a component of the basin of attraction is described in terms of symbolic sequences. We do not give a precise symbolic characterisation of the boundary, but provide a necessary and a sufficient conditions, which is enough for our purposes. To fix notation, we consider the component U_0 .

Let

$$\Sigma_{\underline{k}} = \{(s_0, \underline{k}, s_1, \underline{k}, \dots) : s_0, s_1, \dots \in \mathbb{Z}\},$$

$$\Sigma'_{\underline{k}} = \{(s_0, \underline{k}, s_1, \underline{k}, \dots) : s_0, s_1, \dots \in \mathbb{Z} \text{ and } s_n \notin \{k_1, \dots, k_{p-1}\} \text{ for infinitely many } n\}.$$

Proposition 3.

$$\{z \in J(f) : z \text{ has itinerary in } \Sigma'_{\underline{k}}\} \subset \partial U_0 \subset \{z \in J(f) : z \text{ has itinerary in } \Sigma_{\underline{k}}\}.$$

Proof. First, we show $\partial U_0 \subset \{z \in J(f) : z \text{ has itinerary in } \Sigma_{\underline{k}}\}$. Take $z \in \partial U_0$ and $n \geq 0$. We need to check that $f^n(z) \in \mathcal{H}_{k_j}$ if $n = lp + j$ for an integer l and $j \in \{1, \dots, p-1\}$. It is clear that $f^{lp+j}(z) \in \partial U_j$. Moreover, $\overline{U_j}$ must be contained in $\mathcal{H}_{k_j}$, since it is a connected subset of $\bigcup_s \mathcal{H}_s$ containing $z_j \in C_j \subset \mathcal{H}_{k_j}$. This shows $f^{lp+j}(z) \in \mathcal{H}_{k_j}$.

Now we show $\{z \in J(f) : z \text{ has itinerary in } \Sigma'_{\underline{k}}\} \subset \partial U_0$. Take a point $z \in J(f)$ with itinerary in $\Sigma'_{\underline{k}}$. By definition, there are arbitrarily large n such that $f^{np}(z) \in \mathcal{H}_{s_n}$ and $s_n \neq k_j$ for $j \in \{1, \dots, p-1\}$. This implies that for these n we have $\mathcal{H}_{s_n} \subset \mathcal{L}$. Let I_n be a open vertical segment in $\mathcal{H}_{s_n}$ containing $f^{np}(z)$, such that both endpoints of I_n are in $\partial \mathcal{H}_{s_n}$. Note that the length of I_n is less than 2π because of (1). Let G^n be the branch of f^{-np} along the trajectory of z , i.e. $G^n = G_{s_0, \underline{k}} \circ \dots \circ G_{s_{n-1}, \underline{k}}$. Since $I_n \subset \mathcal{H}_{s_n} \subset \mathcal{L}$, the branch G^n is defined on I_n . By Lemma 2, the length of $G^n(I_n)$ tends to 0 as $n \rightarrow \infty$. Moreover, by the definition of the sets C_j , each map $G_{s_l, \underline{k}}$ maps points from U_0 to points from U_0 . Hence, the endpoints of the curve $G^n(I_n)$ are in U_0 . Since $z \in G^n(I_n)$, this shows $z \in \partial U_0$. $\square$

3. ESCAPING POINTS IN THE BOUNDARY

In this section we prove:

Proposition 4. *The Hausdorff dimension of the set*

$$\{z \in J(f) : z \text{ has itinerary in } \Sigma_{\underline{k}}\} \cap I(f)$$

is equal to 1.

In the proof of this proposition, it is more convenient to consider the set

$$\tilde{\Sigma}_{\underline{k}} = \{(\underline{k}, s_1, \underline{k}, s_2, \dots) : s_1, s_2, \dots \in \mathbb{Z}\}$$

instead of $\Sigma_{\underline{k}}$. Note that the set $\{z \in J(f) : z \text{ has itinerary in } \Sigma_{\underline{k}}\} \cap I(f)$ is a countable union of the images of $\{z \in J(f) : z \text{ has itinerary in } \tilde{\Sigma}_{\underline{k}}\} \cap I(f)$ under inverse branches of f , so both sets have the same Hausdorff dimension. Hence, to prove Proposition 4, it is sufficient to show that

$$(3) \quad \dim_H(\{z \in J(f) : z \text{ has itinerary in } \tilde{\Sigma}_{\underline{k}}\} \cap I(f)) = 1.$$

Let

$$W_M = \{z : \operatorname{Re}(z) \geq M\}$$

for a large integer $M > 0$ and

$$A_M = \{z \in J(f) : z \text{ has itinerary in } \tilde{\Sigma}_{\underline{k}} \text{ and } f^n(z) \in W_M \text{ for all } n \geq 0\}.$$

The assertion (3) will follow immediately from

Proposition 5. *For every $\delta > 0$ there exists $M(\delta) > 0$ such that for every $M > M(\delta)$,*

$$\dim_H A_M < 1 + \delta.$$

The rest of this section is devoted to the proof of Proposition 5.

Definition. Let

$$Q_s^r = \mathcal{L} \cap \mathcal{H}_s \cap \{z : r \leq \operatorname{Re}(z) < r + 1\}$$

for $r, s \in \mathbb{Z}$, $r \geq M$ and let $\mathcal{Q}^r$ be the family of all connected components of $Q_{k_1}^r$.

Remark 6. Note that if M is sufficiently large, then by Lemma 1, each family $\mathcal{Q}^r$ consists of at most p sets.

Let

$$\mathcal{K}_0 = \bigcup_{r=M}^{\infty} \mathcal{Q}^r$$

and for $n \geq 0$ define inductively

$$\mathcal{K}_{n+1} = \{G_{\underline{k},s}(K) : K \in \mathcal{K}_n, s \in \mathbb{Z} \text{ and } G_{\underline{k},s}(K) \cap W_M \neq \emptyset\}.$$

Note that if $x \in A_M$, then for every $n \geq 0$ we have $f^{np}(x) \in \mathcal{L} \cap \mathcal{H}_{k_1} \cap W_M$, so $f^{np}(x)$ belongs to some set $K_0 \in \mathcal{K}_0$. Since

$$\mathcal{L} \cap \mathcal{H}_{k_1} \cap W_M = \bigcup_{K_0 \in \mathcal{K}_0} K_0,$$

this implies that the collection $\mathcal{K}_n$ is a cover of A_M .

Fix $\delta > 0$. We will show that

$$(4) \quad \sup\{\operatorname{diam} K : K \in \mathcal{K}_n\} \rightarrow 0 \text{ as } n \rightarrow \infty.$$

and

$$(5) \quad \sum_{K \in \mathcal{K}_n} (\operatorname{diam} K)^{1+\delta} < 1$$

for every $n \geq 0$, which will prove Proposition 5 by the definition of the Hausdorff measure.

To show (4) and (5), we first prove two simple lemmas.

Lemma 7. *For every sufficiently large M there exists $M_1 > 0$, such that $M_1 \rightarrow \infty$ as $M \rightarrow \infty$, and for every $z \in \mathcal{L} \cap W_M$ and $s \in \mathbb{Z}$,*

$$f^j(G_{\underline{k},s}(z)) \in W_{M_1}$$

for every $j = 1, \dots, p-1$.

Proof. It is sufficient to notice that $\operatorname{Re}(g_s(z)) = \ln|z| - \ln|\lambda| \geq \ln(\operatorname{Re}(z)) - \ln|\lambda|$. $\square$

Lemma 8. *For every $z \in \mathcal{L} \cap W_M$ for sufficiently large M ,*

$$|G'_{\underline{k},s}(z)| < \frac{1}{\operatorname{Re}(z)(\pi|s| + 1)}.$$

Proof. By the chain rule,

$$|G'_{\underline{k},s}(z)| = |g'_s(z)| |g'_{k_{p-1}}(g_s(z))| |(g_{k_1} \circ \cdots \circ g_{k_{p-2}})'(g_{k_{p-1}} \circ g_s(z))|.$$

We have $|g'_s(z)| = 1/|z| \leq 1/\operatorname{Re}(z)$. Moreover, by Lemma 1, there exists a constant $c > 0$, such that

$$|g'_{k_{p-1}}(g_s(z))| = \frac{1}{|g_s(z)|} \leq \frac{2}{(|\operatorname{Re}(g_s(z))| + |\operatorname{Im}(g_s(z))|)} < \frac{2}{\ln M + 2\pi|s| - c} < \frac{1}{\pi|s| + 1}$$

for large M . Finally, by Lemma 7,

$$|(g_{k_1} \circ \cdots \circ g_{k_{p-2}})'(g_{k_{p-1}} \circ g_s(z))| = \prod_{j=1}^{p-2} \frac{1}{|f^j(G_{\underline{k},s}(z))|} \leq \prod_{j=1}^{p-2} \frac{1}{\operatorname{Re}(f^j(G_{\underline{k},s}(z)))} < \frac{1}{M_1^{p-2}} \leq 1,$$

if M is sufficiently large. $\square$

Note that by Lemma 1, for every $K_0 \in \mathcal{K}_0$ we have

$$(6) \quad \operatorname{diam} K_0 < \sqrt{1 + 16\pi^2}.$$

Moreover, by Lemma 8, for every $K \in \mathcal{K}_n$,

$$\operatorname{diam} G_{\underline{k},s}(K) < \frac{\operatorname{diam} K}{M - \operatorname{diam} K},$$

which shows (4) by a simple induction. Moreover, (4) and Lemma 8 imply immediately the following fact.

Corollary 9. *For every $K \in \mathcal{K}_n$ and every $z \in K$,*

$$|G'_{\underline{k},s}(z)| < \frac{1}{(M - d)(\pi|s| + 1)}$$

for some constant $d > 0$. $\square$

Now we are ready to prove (5). Using Remark 6, Lemma 8 and Corollary 9 we get, for some constants $c_1, c_2 > 0$,

$$\begin{aligned}
 \sum_{K \in \mathcal{K}_n} (\text{diam } K)^{1+\delta} &= \sum_{K_0 \in \mathcal{K}_0} \sum_{\substack{s_1, \dots, s_n \in \mathbb{Z}: \\ G_{\underline{k}, s_j} \circ \dots \circ G_{\underline{k}, s_n}(K_0) \in \mathcal{K}_{n-j+1} \\ \text{for } j=1, \dots, n}} (\text{diam } G_{\underline{k}, s_1} \circ \dots \circ G_{\underline{k}, s_n}(K_0))^{1+\delta} \\
 &\leq \sum_{r=M}^{\infty} \sum_{K_0 \in \mathcal{Q}^r} \sum_{\substack{s_1, \dots, s_n \in \mathbb{Z}: \\ G_{\underline{k}, s_j} \circ \dots \circ G_{\underline{k}, s_n}(K_0) \in \mathcal{K}_{n-j+1} \\ \text{for } j=1, \dots, n}} \sup_{z \in K_0} \left(|G'_{\underline{k}, s_n}(z)| \prod_{j=1}^{n-1} |G'_{\underline{k}, s_j}(G_{\underline{k}, s_{j+1}} \circ \dots \circ G_{\underline{k}, s_n}(z))| \right)^{1+\delta} \\
 &< p \sum_{r=M}^{\infty} \sum_{s_1, \dots, s_n \in \mathbb{Z}} \left(\frac{1}{r(\pi|s_n|+1)} \prod_{j=1}^{n-1} \frac{1}{(M-d)(\pi|s_j|+1)} \right)^{1+\delta} \\
 &\leq p \sum_{r=M}^{\infty} \frac{1}{r^{1+\delta}} \left(\frac{1}{(M-d)^{1+\delta}} \right)^{n-1} \left(\sum_{s \in \mathbb{Z}} \frac{1}{(\pi|s|+1)^{1+\delta}} \right)^n < \frac{pc_1}{M^\delta} \frac{c_2^n}{(M-d)^{(1+\delta)(n-1)}} < 1
 \end{aligned}$$

if M is sufficiently large, which shows (5) and ends the proof of Proposition 5.

4. NON-ESCAPING POINTS IN THE BOUNDARY

Proposition 10. *The Hausdorff dimension of the set*

$$\{z \in J(f) : z \text{ has itinerary in } \Sigma'_{\underline{k}} \text{ and the forward orbit of } z \text{ under } f \text{ is bounded}\}$$

is greater than 1.

Proof. The basic idea of the proof is to construct a finite conformal Iterated Function System with a compact Cantor repeller X contained in the considered set, such that $f^{2p}(X) = X$, and to use classical Bowen's formula to estimate the Hausdorff dimension of X . In fact, it is a modification of the proof from [3]. For completeness, we present the main ideas of the proof.

Let R be a large positive real number. Consider the square

$$Q = \left[\frac{R}{2}, \frac{3R}{2} \right] \times \left[\frac{R}{2}, \frac{3R}{2} \right].$$

Note that $Q \subset \mathcal{L}$ for sufficiently large R , so all inverse branches of the form $G_{u, \underline{k}} \circ G_{s, \underline{k}}$, $u, s \in \mathbb{Z}$ are defined on Q . Moreover, $\text{diam } Q = \sqrt{2}R$ and $\text{dist}(Q, \partial \mathcal{L}) > R/2 - D$ for some constant $D > 0$. Hence, by the Koebe Lemma (see e.g. [7]), the distortion of these branches on Q is bounded by a constant independent of R, u, s .

Let

$$Q_{u,s} = G_{u, \underline{k}} \circ G_{s, \underline{k}}(Q)$$

for $u, s \in \mathbb{Z}$ and

$$\mathcal{G} = \{(u, s) \in \mathbb{Z}^2 : Q_{u,s} \subset Q\}.$$

Now define

$$X = \bigcap_{n=1}^{\infty} \bigcup_{i_1, \dots, i_n} h_{i_1} \circ \dots \circ h_{i_n}(Q),$$

where $h_{ij} = G_{u, \underline{k}} \circ G_{s, \underline{k}}$ for some $(u, s) \in \mathcal{G}$. By construction,

$X \subset \{z \in J(f) : z \text{ has itinerary in } \Sigma'_{\underline{k}} \text{ and the forward orbit of } z \text{ under } f \text{ is bounded}\}$
and $f^{2p}(X) = X$.

To prove Proposition 10, we will show that

$$(7) \quad \dim_{\text{H}}(X) > 1.$$

By Bowen's formula (see e.g. [6, 12]), the Hausdorff dimension of X is determined as the unique $t \in \mathbb{R}$, such that $P(t) = 0$ where $P(t)$ is the pressure function:

$$P(t) = \lim_{n \rightarrow \infty} \frac{1}{n} \ln \sum_{i_1, \dots, i_n} \|(h_{i_1} \circ \dots \circ h_{i_n})'\|^t.$$

The function $t \mapsto P(t)$ is strictly decreasing, so to prove (7) it is enough to show that $P(1) > 0$. We have

$$P(1) \geq \lim_{n \rightarrow \infty} \frac{1}{n} \ln \left(\inf_{z \in Q} \sum_{(u, s) \in \mathcal{G}} |(G_{u, \underline{k}} \circ G_{s, \underline{k}})'(z)| \right)^n.$$

Hence, using bounded distortion, to prove (7) it is enough to have

$$(8) \quad \sum_{(u, s) \in \mathcal{G}} |(G_{u, \underline{k}} \circ G_{s, \underline{k}})'(z_R)| > C$$

for a large constant $C > 0$, where z_R is the centre of the square Q , i.e. $z_R = R + iR$.

Now we prove (8). By Lemma 1, for every $z \in \mathcal{H}_s$, $s \in \mathbb{Z}$ and $j = 1, \dots, p-1$, we have

$$(9) \quad |g_{k_j}(z)| \leq |\operatorname{Re}(g_{k_j}(z))| + |\operatorname{Im}(g_{k_j}(z))| \\ < \ln |z| - \ln |\lambda| + (2|s| + 1)\pi + |\operatorname{Arg}(\lambda)| < \ln |z| + 2\pi|s| + c_3$$

for some constant $c_3 > 0$. Using this inductively along the backward trajectory of z_R , we get

$$(10) \quad |G'_{s, \underline{k}}(z_R)| = \frac{1}{|z_R| |g_{k_{p-1}}(z_R)| \cdots |g_{k_1} \circ \dots \circ g_{k_{p-1}}(z_R)|} > \frac{1}{\sqrt{2}R(\ln R + c_4)^{p-1}}$$

and

$$(11) \quad \operatorname{Re}(G_{s, \underline{k}}(z_R)) \leq |G_{s, \underline{k}}(z_R)| < \ln R + c_4$$

for some constant $c_4 > 0$. Note that the points

$$v_s = G_{s, \underline{k}}(z_R)$$

for $s \in \mathbb{Z}$ are located on the same vertical line

$$\ell = \{z : \operatorname{Re}(z) = R_1\}.$$

Let

$$\ell' = \ell \cap \{z : \operatorname{Im}(z) \geq R\}.$$

In the same way as in the proof of Lemma 7, it is easy to see that R_1 is arbitrarily large for large R , so $\ell' \subset \mathcal{L}$ and

$$(12) \quad \operatorname{dist}(z, \partial\mathcal{L}) > D \quad \text{for every } z \in \ell'.$$

For $u \in \mathbb{Z}$ the set $G_{u,\underline{k}}(\ell')$ is a curve tending to infinity and contained in $\mathcal{H}_u$. Moreover, using (11) and (9) inductively along the backward trajectory of ℓ' , we get

$$\inf_{z \in G_{u,\underline{k}}(\ell')} \operatorname{Re}(z) < \ln R + c_4 < \frac{R}{2},$$

if R is sufficiently large. Hence, each $G_{u,\underline{k}}(\ell')$ intersects the strip $\{z : R/2 + 1 < \operatorname{Re}(z) < 3R/2 - 1\}$ along a curve ℓ'_u of length at least $R - 2$. By Lemma 1, there are at least $c_5 R$ such curves ℓ'_u contained in

$$Q' = \left[\frac{R}{2} + 1, \frac{3R}{2} - 1 \right] \times \left[\frac{R}{2} + 1, \frac{3R}{2} - 1 \right],$$

for some constant $c_5 > 0$. By (12) and the Koebe Lemma, the distortion of $G_{u,\underline{k}}$ is uniformly bounded on every vertical segment of endpoints v_s, v_{s+1} , so

$$(13) \quad R - 2 \leq \operatorname{length}(\ell'_u) < c_6 \sum_{s: G_{u,\underline{k}}(v_s) \in \ell'_u} |G'_{u,\underline{k}}(v_s)|$$

for some constant $c_6 > 0$. Moreover, $\operatorname{diam} Q = \sqrt{2}R$ and $(G_{u,\underline{k}} \circ G_{s,\underline{k}})'(z) < 1/(\sqrt{2}R)$ for $z \in Q$ (the proof is analogous to the proof of Lemma 8), so

$$\operatorname{diam} Q_{u,s} < 1.$$

Hence, if $G_{u,\underline{k}}(v_s) \in \ell'_u \subset Q'$, then $Q_{u,s} \subset Q$, so $(u, s) \in \mathcal{G}$. Using (13), we conclude that

$$\sum_{(u,s) \in \mathcal{G}} |G'_{u,\underline{k}}(v_s)| \geq \sum_{u: \ell'_u \subset Q'} \sum_{s: G_{u,\underline{k}}(v_s) \in \ell'_u} |G'_{u,\underline{k}}(v_s)| > \frac{c_5}{c_6} R(R - 2).$$

Combining this with (10), we get

$$\sum_{(u,s) \in \mathcal{G}} |(G_{u,\underline{k}} \circ G_{s,\underline{k}})'(z_R)| = \sum_{(u,s) \in \mathcal{G}} |G'_{u,\underline{k}}(v_s)| |G'_{s,\underline{k}}(z_R)| > \frac{c_5}{\sqrt{2}c_6} \frac{R - 2}{(\ln R + c_4)^{p-1}},$$

which shows (8) for sufficiently large R . This completes the proof. $\square$

Note that it follows from the above proof that the set $X \cup f^p(X)$ is a conformal f^p -invariant Cantor set and $|(f^p)'| > 1$ on $X \cup f^p(X)$. Hence we immediately obtain the following.

Proposition 11. *The hyperbolic dimension of the set*

$$\{z \in J(f) : z \text{ has itinerary in } \Sigma'_{\underline{k}} \text{ and the forward orbit of } z \text{ under } f \text{ is bounded}\}$$

is greater than 1. $\square$

5. CONCLUSION

By Propositions 3, 4 and 10, the set $\partial U_0 \cap I(f)$ has Hausdorff dimension 1, while $\partial U_0 \cap J_{\text{bd}}(f)$ has Hausdorff dimension greater than 1. This implies

$$1 < \dim_{\text{H}}(\partial U_0 \cap J_{\text{bd}}(f)) \leq \dim_{\text{H}}(\partial U_0 \setminus I(f)) = \dim_{\text{H}}(\partial U_0).$$

Of course, the same holds for any component U of the entire basin of attraction. This proves the first part of Theorem A, the lower estimate of the first part and the second part of Theorem B. The remaining part is a consequence of the second part of Theorem B and the fact $\dim_{\text{H}}(J(f) \setminus I(f)) < 2$, which was proved in [15] as Theorem 6.1. Finally, the remark after Theorem B follows from Propositions 3 and 11.

REFERENCES

- [1] J. M. Aarts and L. G. Oversteegen, *The geometry of Julia sets*, Trans. Amer. Math. Soc. 338 (1993), 897–918.
- [2] I. N. Baker, *The domains of normality of an entire function*, Ann. Acad. Sci. Fenn. Ser. A I Math. (1975), 1, 277–283.
- [3] K. Barański, B. Karpińska and A. Zdunik, *Hyperbolic dimension of Julia sets of meromorphic maps with logarithmic tracts*, Internat. Math. Res. Notices (2008), Vol. 2008, Article ID rnn141, 10 pages.
- [4] R. Bhattacharjee and R. L. Devaney, *Tying hairs for structurally stable exponentials*, Ergodic Theory Dyn. Syst. 20 (2000), 1603–1617.
- [5] R. Bhattacharjee, R. L. Devaney, R. E. Lee Deville, K. Josić and M. Moreno-Rocha, *Accessible points in the Julia sets of stable exponentials*, Discrete Contin. Dyn. Syst., Ser. B 1 (2001), No. 3, 299–318.
- [6] R. Bowen, *Hausdorff dimension of quasicircles*, Inst. Hautes Études Sci. Publ. Math. 50 (1979), 11–25.
- [7] L. Carleson and T. W. Gamelin, *Complex dynamics*, Springer-Verlag, New York, 1993.
- [8] B. Karpińska, *Area and Hausdorff dimension of the set of accessible points of the Julia sets of λe^z and $\lambda \sin z$* , Fund. Math. 159 (1999), 269–287.
- [9] B. Karpińska, *Hausdorff dimension of the hairs without endpoints for $\lambda \exp z$* , C. R. Acad. Sci. Paris Ser. I Math. 328 (1999), 1039–1044.
- [10] B. Karpińska and M. Urbański, *How points escape to infinity under exponential maps*, J. London Math. Soc. 73 (2006), 141–156.
- [11] C. T. McMullen, *Area and Hausdorff dimension of Julia sets of entire functions*, Trans. Amer. Math. Soc. 300 (1987), 329–342.
- [12] F. Przytycki and M. Urbański, *Fractals in the Plane – the Ergodic Theory Methods*, to appear in Cambridge University Press, available on-line at www.math.unt.edu/~urbanski.
- [13] D. Schleicher and J. Zimmer, *Escaping points of exponential maps*, J. London Math. Soc. 67 (2003), 380–400.
- [14] M. Urbański and A. Zdunik, *Real analyticity of Hausdorff dimension of finer Julia sets of exponential family*, Ergodic Theory Dynam. Systems 24 (2004), 279–315.
- [15] M. Urbański and A. Zdunik, *The finer geometry and dynamics of the hyperbolic exponential family*, Michigan Math. J. 51 (2003), 227–250.

INSTITUTE OF MATHEMATICS, UNIVERSITY OF WARSAW, UL. BANACHA 2, 02-097 WARSZAWA,
POLAND

E-mail address: `baranski@mimuw.edu.pl`

FACULTY OF MATHEMATICS AND INFORMATION SCIENCE, WARSAW UNIVERSITY OF TECHNOLOGY,
PL. POLITECHNIKI 1, 00-661 WARSZAWA, POLAND

E-mail address: `bkarpin@mini.pw.edu.pl`

INSTITUTE OF MATHEMATICS, UNIVERSITY OF WARSAW, UL. BANACHA 2, 02-097 WARSZAWA,
POLAND

E-mail address: `A.Zdunik@mimuw.edu.pl`